



**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In re Patent Application of

MAVILIO

Atty. Ref.: 1303-110; Confirmation No. 5693

Appl. No. 09/674,853

TC/A.U. 1632

Filed: November 7, 2000

Examiner: Wehbe

For: GENETICALLY-MODIFIED FIBROBLASTS  
AND THE USE THEREOF

\* \* \* \* \*

July 14, 2004

**MAIL STOP AF**

Commissioner for Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

Sir:

**SUBMISSION**

Submitted herewith is an English translation of the priority document MI98 A  
001004.

Attached is a Notice of Appeal to maintain pendency.

A favorable response to the Amendment of May 14, 2004 is earnestly solicited  
without further delay. The USPTO PAIR page confirms the applicants Amendment of  
Mail 14, 2004 has been received. See, attached print-out of PAIR page. A response to  
the Amendment however has yet to be received by the undersigned. The attached  
appears to confirm that the Examiner has not yet issued a response to the Amendment  
of May 14, 2004.


MAVILIO  
Appl. No. 09/674,853  
July 14, 2004

The Examiner is requested to contact the undersigned in the event anything further is required to place the application in condition for allowance.

Respectfully submitted,

**NIXON & VANDERHYE P.C.**

By: \_\_\_\_\_




B. J. Sadoff  
Reg. No. 36,663

BJS  
1100 North Glebe Road, 8th Floor  
Arlington, VA 22201-4714  
Telephone: (703) 816-4000  
Facsimile: (703) 816-4100



## United States Patent and Trademark Office

[Home](#) | [Site Index](#) | [Search](#) | [Guides](#) | [Contacts](#) | [eBusiness](#) | [eBiz alerts](#) | [News](#) | [Help](#)
**PATENT APPLICATION INFORMATION RETRIEVAL**

Search results for application number: 09/674,853			
Application Number:	09/674,853	Customer Number:	23117
Filing or 371(c) Date:	11-07-2000	Status:	Final Rejection Mailed
Application Type:	Utility	Status Date:	01-14-2004
Examiner Name:	WEHBE, ANNE MARIE SABRINA	Location:	ELECTRONIC
Group Art Unit:	1632	Location Date:	-
Confirmation Number:	5693	Earliest Publication No:	-
Attorney Docket Number:	1303-110	Earliest Publication Date:	-
Class/ Sub-Class:	424/093.21	Patent Number:	-
First Named Inventor:	Fulvio Mavilio, Milano, (IT)	Issue Date of Patent:	-
Title Of Invention:	Genetically-modified fibroblasts and the use thereof		

## Search Options

Assignments
Foreign Priority
Image File Wrapper
Publication Review

File History	
Date	Contents Description
05-14-2004	Workflow incoming amendment IFW
01-14-2004	Mail Final Rejection (PTOL - 326)
01-12-2004	Final Rejection
10-30-2003	Date Forwarded to Examiner
10-17-2003	Response after Non-Final Action
10-17-2003	Request for Extension of Time - Granted
06-17-2003	Mail Non-Final Rejection
06-16-2003	Non-Final Rejection
04-02-2003	Date Forwarded to Examiner
03-26-2003	Request for Continued Examination (RCE)
04-02-2003	Express Abandonment (for Entry of CPA / RCE / Rule129)
03-26-2003	Workflow - Request for RCE - Begin
01-31-2003	Mail Advisory Action (PTOL - 303)
01-31-2003	Advisory Action (PTOL-303)

01-02-2003	Correspondence Address Change
01-02-2003	Date Forwarded to Examiner
12-26-2002	Amendment after Final Rejection
09-26-2002	Mail Final Rejection (PTOL - 326)
09-26-2002	Final Rejection
07-10-2002	X-Pre-Legal Complete Amended Case
07-10-2002	Date Forwarded to Examiner
07-01-2002	Response after Non-Final Action
03-29-2002	Mail Non-Final Rejection
03-26-2002	X-Post-Legal Complete Rejection
03-25-2002	Non-Final Rejection
01-31-2001	Case Docketed to Examiner in GAU
01-22-2001	Application Dispatched from OIPE
01-18-2001	IFW Scan & PACR Auto Security Review
01-09-2001	Correspondence Address Change
12-08-2000	Released to OIPE
12-08-2000	Notice of DO/EO Acceptance Mailed
12-07-2000	371 Application Preexamination Docketing
11-16-2000	371 Application Preexamination Docketing
11-07-2000	Receipt of 371 Request
11-16-2000	Correspondence Address Change
11-07-2000	Initial Exam Team nn




United States Patent and Trademark Office

[Home](#) | [Site Index](#) | [Search](#) | [Guides](#) | [Contacts](#) | [eBusiness](#) | [eBiz alerts](#) | [News](#) | [Help](#)



## PATENT APPLICATION INFORMATION RETRIEVAL



### Other Links




Image File Wrapper for Application No.:09/674,853						
Mail Room Date	Document Description	Document Category	Page Count	All	None	Download
05/14/2004	<u>Claim</u>	PROSECUTION	2		<input type="checkbox"/>	
05/14/2004	<u>Applicant Arguments or Remarks Made in an Amendment</u>	PROSECUTION	4		<input type="checkbox"/>	
05/14/2004	<u>Miscellaneous Incoming Letter</u>	PROSECUTION	1		<input type="checkbox"/>	
05/14/2004	<u>Amendment After Final</u>	PROSECUTION	1		<input type="checkbox"/>	
01/14/2004	<u>NPL Documents</u>	PROSECUTION	7		<input type="checkbox"/>	
01/14/2004	<u>List of references cited by Examiner</u>	PROSECUTION	1		<input type="checkbox"/>	
01/14/2004	<u>Search info on File Wrapper</u>	PROSECUTION	1		<input type="checkbox"/>	
01/14/2004	<u>Final Rejection</u>	PROSECUTION	15		<input type="checkbox"/>	
10/17/2003	<u>Amendment</u>	PROSECUTION	1		<input type="checkbox"/>	
10/17/2003	<u>Applicant Arguments or Remarks Made in an Amendment</u>	PROSECUTION	11		<input type="checkbox"/>	
10/17/2003	<u>Claim</u>	PROSECUTION	3		<input type="checkbox"/>	
06/17/2003	<u>List of references cited by Examiner</u>	PROSECUTION	1		<input type="checkbox"/>	
06/17/2003	<u>Non-Final Rejection</u>	PROSECUTION	12		<input type="checkbox"/>	
03/26/2003	<u>Request for Continued Examination (RCE)</u>	PROSECUTION	1		<input type="checkbox"/>	
01/31/2003	<u>Advisory Action (PTOL-303)</u>	PROSECUTION	4		<input type="checkbox"/>	
12/26/2002	<u>Change of Address</u>	PROSECUTION	1			

[http://pair-direct.uspto.gov/cgi-bin/final/pair/pair\\_efw.pl](http://pair-direct.uspto.gov/cgi-bin/final/pair/pair_efw.pl)

**Sorted By : Document Category**

[|.HOME](#) | [INDEX](#) | [SEARCH](#) | [eBUSINESS](#) | [CONTACT US](#) | [PRIVACY STATEMENT](#)



**MINISTRY OF INDUSTRY, TRADE AND HANDICRAFT  
GENERAL DIRECTORATE OF INDUSTRIAL PRODUCTION  
CENTRAL PATENT OFFICE**

ITL. 20,000 revenue stamps and seal of the  
Ministry of Industry, Trade and Handicraft.  
Central Patent Office.

**Certification of copy of documents relating to a patent application for industrial  
invention N. MI98 A 001004**

It is hereby declared that the enclosed copy  
corresponds to the documents as originally filed  
with the above mentioned patent application whose  
data are as from the enclosed filing certificate.

Rome, May 18, 1999

By proxy of  
THE DIRECTOR OF THE DIVISION  
Dr. Paola DI CINTIO  
(signature)

WHITE, RED AND GREEN RIBBON  
AND ORANGE SEAL OF THE MINI-  
STRY OF INDUSTRY, TRADE AND  
HANDICRAFT.  
CENTRAL PATENT OFFICE

**TO THE MINISTRY OF INDUSTRY, TRADE AND**

Space to be covered  
with stamp

**HANDICRAFT**

**CENTRAL PATENT OFFICE - ROME**

**FORM A**

APPLICATION FOR INDUSTRIAL INVENTION,  
FILING MISSING DOCUMENTS, ADVANCED A-  
VAILABILITY TO THE PUBLIC

**A. APPLICANT(S)**

NAME **FONDAZIONE CENTRO SAN RAFFAELE DEL MONTE TABOR** EN  
1) RESIDING IN **Milan** CODE **03064280153**  
2) NAME [ ]  
RESIDING IN CODE

**B. APPLICANT'S REPRESENTATIVE AT THE CENTRAL PATENT OFFICE**

SURNAME, NAME: **BIANCHETTI Giuseppe et al** FISCAL CODE:

NAME OF THE OFFICE: **Bianchetti • Bracco • Minoja s.r.l.**

STREET: **Rossini N. 8** TOWN: **MILAN** ZIP CODE: **20122** PROV.: **MI**

**C. ADDRESSEE'S ELECTIVE DOMICILE:**

STREET: N. TOWN: ZIP CODE: PROV:

**D. TITLE** proposed class(sect/cl/scl):[**C12N**] Group/sub-group:[**15/00**]

**"Genetically modified fibroblasts and the use thereof"**

**ADVANCED ACCESS TO THE PUBLIC:** YES[ ] NO[X]

IF REQUEST: DATE / / RECORD N. [ ]

**E. NAMED INVENTORS:**

	surname, name	surname, name
1)	<b>Mavilio Fulvio</b>	3)
2)		4)

**F. PRIORITY**

COUNTRY OR ORGANIZATION	KIND OF PRIORITY	APPL.N.	FILING DATE	ENCL Y/M
1)			/ /	
2)			/ /	

FILING DATE

/ /

/ /

**ENCLOSED DOCUMENTS:**

N. COPIES

//

//

//

//

///

**Five hundred sixty-five thousand=**  
compulsory

**SIGNATURE OF THE APPLICANT(S)**

**Minoja Fabrizio (signature)**

CODE [15]

APPLICATION NUMBER MI98A 001004 Reg.A

**I. VARIOUS NOTES OF THE ATTESTING OFFICER**

**THE PETITIONER**

Biglia Michele  
(signature)

**SEAL OF THE OFFICE**

Seal of the Ministry  
of Industry, Commerce  
and Handicraft  
ITL. 20,000.- revenue  
stamps.

**THE ATTESTING OFFICER**

CORTONESI MAURIZIO  
(signature)

**ITL. 20,000.- revenue stamps  
Seal of the Ministry of Industry,  
Commerce and Handicraft**

# ABSTRACT OF THE INVENTION WITH MAIN DRAWING, DESCRIPTION AND CLAIMS

Application N. **MI98 A 001004** Reg.A Filing date **08/05/1998**  
 Patent N. Granting date / /

#### D. TITLE

## **"Genetically modified fibroblasts and the use thereof"**

## L. ABSTRACT

A method for the preparation of genetically-modified fibroblasts expressing a muscle lineage commitment gene, and the use thereof for the treatment of genetic defects or for the expression of therapeutic proteins.

## M. DRAWING

Description of the industrial invention of title:

**"GENETICALLY-MODIFIED FIBROBLASTS AND THE USE THEREOF"**

in the name of: FONDAZIONE CENTRO SAN RAFFAELE DEL MONTE TABOR  
residing in: Milan

\* \* \*

The present invention relates to an ex-vivo method for the myogenic conversion of genetically-modified fibroblasts, for use in the correction of congenital defects of the muscular system, such as primary monogenic myopathies, or in the production of therapeutical proteins.

5 Furthermore, the invention relates to the use of genetically-modified fibroblasts for the preparation of compositions for cell implants for the treatment of muscle pathologies with a genetic aetiology, or for the in vivo secretion of recombinant proteins of therapeutical importance. In particular, the invention is valuable for the treatment of muscular  
10 dystrophies, a heterogeneous group of severe degenerative diseases of the muscle caused by mutations in the genes encoding for the membrane-associated protein dystrophin or for other members of the dystrophin-associated protein complex connecting the muscle fiber cytoskeleton with the extracellular matrix.

15 Until now, the attempts to correct muscular dystrophy with gene therapy based on the implantation of muscles or other tissues stem cells, have been thwarted by enormous difficulties in the isolation of a suitable number of myogenic cells to be genetically-modified or to the poor yield of phenotypic conversion. By way of example, the difficulty in obtaining  
20 a reasonable number of satellite cells to be genetically-modified is a limit to gene therapy of patients with Duchenne muscular dystrophy or with Becker muscular dystrophy (Salvatori, G., et al., 1993; Webster, C. et al., 1990). Further problems related to the implantation of genetically-modified cells in dystrophic patients are the immunogenicity of viral

vectors, the difficulty in developing an effective administration route and the poor survival of injected cells. Furthermore, the expression of dystrophin itself has been reported to induce cellular and/or humoral immune responses in dystrophin-lacking patients (New England J. of Medicine 333, 732-733, 1995).

It has been known for some time that forced expression of MyoD, and related members of the myogenic family of basic helix-loop-helix transcription factors, would activate myogenesis in non-muscle cells (Davis, R. L., et al., 1987; Cossu, G., et al., 1996; Cossu, G., 1997). The Applicant itself has recently reported that fibroblasts undergo myogenic differentiation when co-cultured with myogenic lines (C2C12) or primary cells, but not with other cell types (Breton, M. et al., 1995; Gibson, A.J., et al., 1995; Salvatori, G. et al., 1995). WO 95/12979 discloses a general method for inducing cell differentiation to a novel phenotype, for example muscular phenotype, by inserting the genic sequences of factors such as MyoD, myogenin, Myf-5 and MRF4 into the cells. The same document neither considers any possibility to convert genetically-modified cells to the myogenic phenotype in order to make them suitable for gene therapy, nor envisages the possibility to carry out an "ex vivo" myogenic conversion.

GB 2293604 discloses the use of fibroblasts in the treatment of muscular pathologies, wherein fibroblasts themselves can be transformed with a therapeutically active gene, for example the dystrophin gene, whereas the possibility to induce differentiation by means of myogenic factors has not been considered.

In light of what stated above, it appears remarkably advantageous to rely on a method which allows a high number of cells to be converted

to the myogenic phenotype, after being genetically-modified ex-vivo, in order to correct a genetic defect, such as muscular dystrophy, or any other dysfunction related, for instance, to insufficient or incorrect production of a protein or a plasma factor or other secreted or circulating proteins, as in  
5 the case of diabetes, hemophilia, pituitary dwarfism.

It has now been found that it is possible to convert genetically-modified fibroblasts to the muscular phenotype, with a high-efficiency process, so as to effectively produce therapeutical proteins or proteins able to correct the genetic defects, particularly muscular dystrophy.

10 According to a first aspect, the invention provides a high-efficiency method for the preparation of genetically-modified fibroblasts expressing muscle lineage commitment genes, which comprises:

- a) ex-vivo transduction of fibroblasts with a therapeutic gene or a gene capable of correcting a genetic defect;
- 15 b) transient expression of the muscle lineage commitment gene in fibroblasts transduced as at point (a), through transformation of the cells with a high efficiency DNA transfer method, preferably a viral vector, more preferably selected from baculovirus, adeno-associated viruses and, most preferably, adenovirus, or with other  
20 high efficiency DNA transfer means, such as liposomes, nucleoprotein complexes, polyethylenimines, wherein the gene of the muscle lineage commitment is under the control of a strong promoter, preferably a viral promoter.

"Muscle lineage commitment gene" means any gene capable of  
25 converting fibroblasts to the myogenic phenotype, particularly MyF-5, MRF4, myogenin and, preferably, MyoD genes.

Therapeutical genes which can be transduced into fibroblasts



before the myogenic conversion are, for example, genes involved the various types of muscle dystrophies, such as sarcoglycan, emerin and, preferably, dystrophin genes, or genes coding for circulating plasma proteins, such as coagulation factors or insulin, for hormones, such as growth hormone, or other genes, such as hypoxanthine-guanine phospho-  
5 ribosyl transferase, adenosine deaminase, purine nucleoside phospho-  
rylase, glucocerebrosidase, low density lipoprotein receptor, phenylalanine hydroxylase, arginine succinate synthetase and aryl-sulfatase.

A second aspect of the invention relates to genetically-modified  
10 fibroblasts, transiently expressing a gene of the muscle lineage commitment, preferably MyoD, obtained by the method of the invention.

The conversion of fibroblasts to the myogenic phenotype by transient expression of a muscle lineage commitment gene, proved to be surprisingly more effective than other alternative techniques. In fact, co-  
15 culture of fibroblasts with satellite cells, treatment with dexamethasone and transfection with calcium phosphate, lipofectamine, or electroporation of the same plasmid expressing MyoD, induced myogenic conversion in a percentage ranging from 1% to 14%, in comparison with values higher than 70% in case of infection with a recombinant adenoviral  
20 vector. Furthermore, the efficiency of conversion was found to increase linearly for m.o.i. ("multiplicity of infection") values from 500 to 2000, whereas cytotoxicity at maximum m.o.i. values, remained within acceptable limits. Similar results were obtained with human or murine fibroblasts, whereas as far as tissutal fibroblasts are concerned, dermal  
25 fibroblasts gave better results. One main advantage of the method of the invention is that transient expression of the exogenous muscle lineage-commitment gene activates the corresponding endogenous gene, therefore

irreversibly committing cells to myogenesis, thus rendering constitutive expression of the transgene unnecessary. The actual myogenic conversion of fibroblasts transiently expressing exogenous MyoD has been confirmed by immuno-histochemistry, electron microscopy and gene expression analysis of different muscle-specific markers, such as myosin light and heavy chains, acetylcholine receptor  $\alpha$ -subunit, M creatine kinase, myogenin and MyoD.

The high efficiency of the method allows the preparation of genetically manipulated fibroblasts in which a therapeutic gene, or a gene correcting the genetic defect in the muscle has been inserted, which may be stably converted to the myogenic phenotype. In fact, conversion to the muscle phenotype prevents cell replication and provides stable implants, based on the use of muscle fibers.

Furthermore, the efficiency of the method as well as the stability of the resulting product allow to successfully carry out gene-therapy, in which genetically-modified fibroblasts expressing a gene of the muscle lineage commitment, after appropriate ex-vivo treatment, are injected into muscle tissue where they are capable of regenerating the muscle fibers and express the correct gene.

A number of evidences proved that fibroblasts treated ex-vivo as herein disclosed are able to regenerate muscle fibers in mice, indistinguishable from those originating from primary myogenic (satellite) cells, with a long-lasting effect. Moreover, after treatment, only antibody mediated immune response against proteins of adenoviral infected fibroblasts has been detected. No cell-mediated immune response has been observed at the site of injection of the modified fibroblasts. This is a significant advantage in gene therapy, as one of the major problems

associated to the in vivo use of adenoviral vectors is the induction of significant, T-cell-mediated immune responses against both the viral and transgenic proteins.

According to a further aspect, the invention relates to the use of  
5 fibroblasts obtained by the above described method, for "ex vivo" gene therapy or for the preparation of compositions for stable cell implants based on muscle fibers.

A typical application of the ex-vivo gene therapy comprises, for example, isolating dermal fibroblasts from a patient suffering from a  
10 genetic disease which alters the muscle structure or functionality, such as dystrophy, expanding the cultured cells, transducing the cells ex-vivo with a retroviral vector containing the correct gene, and inducing myogenesis by infection with an adenoviral vector, or with another vector as mentioned above, containing the gene of the muscle lineage  
15 commitment, and re-implanting modified cells into the muscle tissue.

Another typical approach for the preparation of cell implants based on muscle fibers comprises isolating dermal fibroblasts from a patient suffering from a disease characterised by, for example, lack of a plasma protein (such as diabetes, pituitary dwarfism, hemophilia), expanding the  
20 cultured cells, transducing the cells ex-vivo with a retroviral vector containing the therapeutic gene, and inducing myogenesis by infection with an adenoviral vector, or with another vector as mentioned above, containing the muscle lineage commitment gene, encapsulating the modified cells in the most suitable implantation matrix and implanting the  
25 preparation in the preferred body site. In this specific application, induction of myogenesis and subsequent differentiation into muscle tissue is not aimed at the restoration of muscle functionality, but to the stability

of the implant by arrest of cell division.

As further applications of the present invention, its use in zootechnic field can be envisaged, to induce a specific characteristic in animals, for example an increase in weight, by production of a hormone or any other substance by the cells treated ex vivo with the herein disclosed method and then re-implanted into the animal; or the creation or use of animal models of human pathologies for the study of novel therapeutical procedures to be carried out with the method of the invention.

The following examples illustrate the invention in greater details.

#### **Example 1**

##### **Preparation of genetically-modified fibroblasts expressing MyoD.**

Fetal fibroblasts were isolated from skin and skeletal muscle of C3H mouse embryos (15-17 days) or legally aborted human fetuses (8-12 weeks) as described in Salvatori et al., *Human Gene Ther.*, 1993, 4:713-723, or human dermal or muscle fibroblasts were obtained from tissues of patients undergoing post-traumatic surgery. In this case tissues were finely minced with scissors, digested with 2 mg/ml dispase, 0.1 mg/ml collagenase in phosphate buffered saline (PBS) for 45' at 37°C, washed in RPMI medium, and pipetted to obtain a single-cell suspension.

Murine bone marrow fibroblasts transgenic for the lacZ gene with nuclear localization, were obtained by resuspending and plating cells flushed from the long bones of 8-10-wk-old MLC3F/nlacZ mice. Human bone marrow fibroblasts were obtained from healthy donors after removal of non-adherent cells. All cells were grown in RPMI supplemented with 15% Fetal calf serum (FCS), 1% gentamycin and 0.3 mM  $\beta$ -mercaptoethanol (growth medium). Human bone marrow fibroblasts were

supplemented with 2 ng/ml bFGF.

Myogenic differentiation was induced by shifting the cells to RPMI supplemented with 2% horse serum (differentiation medium). Fibroblasts were purified by sub-culture (at least two rounds) in growth medium.

5 Removal of myogenic cells was routinely verified by immunocytochemical staining of a cell aliquot sub-cultured for 5 days in differentiation medium.

For transduction, fibroblasts (from third to tenth passage) were infected with a replication-defective retroviral vector (LBSN) expressing  
10 a cytoplasmic  $\beta$ -galactosidase gene under the LTR promoter (Salvatori, G., et al., 1993).

For the subsequent step (transient expression of MyoD), cells were treated with the Ad5-derived, E1A-deleted adenoviral vector expressing the full-length murine MyoD c-DNA under the transcriptional control of  
15 Rous Sarcoma Virus (RSV) LTR (Murry, C. E., et al., 1996). In some control experiments, cells were transfected by standard precipitation techniques with calcium-phosphate or lipofectamine (Dotap) with 10  $\mu$ g of the PMC11 plasmid, containing the MyoD cDNA under the control of the CMV promoter.

20 Alternatively, cells were electroporated with 6  $\mu$ g of the same plasmid in growth medium at 120 V, 960 mF. After transduction, cells were grown for 24 hrs in growth medium and then either shifted to differentiation medium for 3-4 days or injected in vivo.

In some experiments, cells were pre-labelled with 0.5 mCi/ml of  
25 [14C] thymidine (Amersham) for 24 h and then exposed to the adenoviral vector expressing MyoD and survival was measured by counting residual cpm incorporated.

### **Example 2**

#### **Implantation of MyoD-converted fibroblasts into mouse muscles.**

1 or 2 millions of transduced (or transgenic) human or murine fibroblasts, converted by transient expression of MyoD, were trypsinized, resuspended in 20-50  $\mu$ l of PBS and injected into a single site of the regenerating tibialis anterior muscle of either syngeneic (C3H) or immunodeficient (scid/bg) mice which had received a 30- $\mu$ l injection of 10<sup>-5</sup> M cardiotoxin (Latoxan) 48 hrs earlier.

For some experiments, myogenically converted mouse fibroblasts were labelled with DiI dye 1 hr before injection in vivo (a 0.5% solution of DiI in absolute ethanol was diluted just before use in 0.3 M sucrose to a final concentration of 0.05%).

Mice were sacrificed after various periods of time, and muscles were cryo-sectioned and stained for  $\beta$ -galactosidase activity or processed for immuno-histochemistry. At the time of sacrifice, serum was collected from C3H mice and reacted with adeno-infected or control cells at various dilutions, followed by a fluorescein-conjugated anti-mouse IgG secondary antibody.

### **Example 3**

#### **Immunocytochemistry, electron microscopy and RNA analysis techniques.**

Immunofluorescence analysis was carried out as described (Cusella - De Angelis, M.G., et al., 1994) using the following antibodies: MF20, a monoclonal antibody which recognises all sarcomeric myosins (Bader, D., et al., 1982); a rabbit antiserum against sarcomeric proteins (Tajbakhsh, S., et al., 1994); an anti-MyoD polyclonal antibody (Hasty, P., et al., 1993); BD5, a monoclonal antibody which recognises myosin heavy

chains (Yasin, R., et al., 1977); a rabbit polyclonal antibody against human fetal myosin (Edom, F., et al., 1994); an anti-bromo-deoxyuridine (BrdU) monoclonal antibody; anti-leu, anti-CD4 and anti-Mac3 rat anti-mouse leukocyte antigen antibodies. Briefly, cells cultures or cryostat  
5 sections were fixed with 4% paraformaldehyde for 10 min at 4°C, washed 3X in PBS, and incubated at 4°C with the primary antibodies in PBS + 1% bovine serum albumin (BSA). BrdU labelled cell-cultures were treated for 10 min at rt with 2M HCl and washed 3X with PBS before incubation with the anti-BrdU antibody.

10 After the first incubation, cells or sections were washed 3X with PBS + 1% BSA and incubated for 1 hr at rt with rhodamine-conjugated goat anti-mouse Ig or with a fluorescein-conjugated goat anti rabbit Ig (1:100 dilution).

Cultures or sections were then washed, mounted in 75%  
15 glycerol/PBS (pH 7.5), and observed under a Zeiss Axiophot epifluorescence microscope.

For electron microscopy cells were washed in PBS, fixed in 2% glutaraldehyde in 0.1 M Millonig's buffer (pH 7.4) for 1 hr at 4°C, post-fixed for 1 hr in 1% buffered osmium tetroxide and dehydrated in  
20 alcohol. Cells were treated with propylene oxide, embedded in Epon 812 and cut into ultrathin sections. The ultrathin sections were stained with uranyl acetate and lead citrate, and examined with a Zeiss 109 electron microscope.

For RNA analysis, RNA was extracted by the method of  
25 Chomczynski et al. (Chomczynski, P., et al., 1987), run in 15 µg aliquots on 1% agarose/formaldehyde gels, and transferred by capillary Northern blotting to nylon membranes (Amersham Hybond-N).

Filters were cross-linked for 2 hrs at 80 °C under vacuum, and hybridized to [32P]-labelled probes for MyoD and myogenin (Bober, E., et al., 1991), MLC1F and MCK (Lyons, G. E., et al., 1991), Ach-receptor  $\alpha$ -subunit (25) under standard conditions (26).

5 **Example 4**

**Analysis of in vitro converted fibroblasts expressing the myogenic phenotype.**

In order to compare the adeno-MyoD-induced phenotype with that of primary myogenic cells, a number of muscle-specific markers were  
10 analysed by immuno-cytochemistry, electron microscopy and gene expression analysis, in human and murine converted dermal fibroblasts, and in differentiating satellite cells of comparable age.

All differentiated myotubes derived from fusion of converted murine newborn dermal fibroblasts express fast embryonic myosin heavy  
15 chains, while a fraction of them also express slow myosin heavy chains, while a fraction of them also express slow myosin heavy chains, as observed in myotubes derived from satellite cells.

Similarly, myotubes derived from human fetal dermal fibroblasts show a well organized initial sarcomerogenesis, with aligned sarcomeres  
20 and patterned Z lines, as reported for myotubes derived from primary myogenic cells, which never complete sarcomerogenesis in vitro.

In order to verify whether the introduction of MyoD in primary fibroblasts would also activate transcription of genes responsible for membrane and metabolic muscle-specific functions, the Applicant  
25 measured by Northern blotting the expression of RNAs coding muscle-specific proteins such as the acetylcholine (ACh) receptor  $\alpha$ -subunit, M creatine kinase (MCK), as well as MyoD, myogenin, and myosin light



chain 1 fast (MLC1F) chosen as positive controls.

No detectable level of any of these messages was detected in unconverted fibroblasts; on the other hand, these mRNAs were expressed at comparable levels in myotubes derived from converted fibroblasts and  
5 in myotubes derived from satellite cells of corresponding age.

Thus, under all the parameters analysed in vitro, myotubes originating from converted fibroblasts cannot be distinguished from those derived by primary myogenic cells.

Afterwards, the Applicant investigated whether fibroblasts exposed  
10 to the MyoD adenoviral vector would still maintain the capacity to divide, since MyoD overexpression has an anti-proliferative effect. Murine and human fetal skin fibroblasts were infected with the MyoD adenoviral vector for 3 hrs at an m.o.i. of 2,000 in serum-free medium, cultured for 12 hrs in growth medium containing 10 mM BrdU, fixed at 12, 24, 48 and  
15 72 hrs after infection, and double-stained with anti-MyoD and anti-BrdU antibodies.

24 hrs after exposure to the adenoviral vector, only a few cells expressing MyoD had incorporated BrdU, while no double-positive cells were detected at 48 and 72 hrs. Expression of MyoD therefore blocks cell  
20 division in converted fibroblasts.

### **Example 5**

#### **Myogenic conversion efficiency of human and murine fibroblasts of different origins**

Fibroblasts were isolated, expanded, infected with the MyoD  
25 adenoviral vector for 3 hrs at an m.o.i. of 2,000, cultured for a further 24 hrs and then induced to differentiation as described. Table 1 shows the conversion % values.

The conversion efficiency was measured as percentage of cells which expressed sarcomeric myosin heavy chains, as shown by staining with anti-myosin antibody. (MF20). Values shown are the average of two separate experiments, each performed in triplicate.

Table 1

		Murine		Human	
		Fetal	Adult	Fetal	Adult
10	Derma	70	56	65	42
	Muscle	43	44	40	32
	Bone marrow	ND	6	ND	6

ND: not done.

**Example 6**

**Survival of MyoD-converted fibroblasts and satellite cells (from MLC3F/LacZ mice) after injection into regenerating Tibialis Anterior muscle (TA) of SCID/bg mice.**

20  $\mu$ l containing  $1 \times 10^6$  cells were injected into a single site of regenerating TA of SCID/bg mice.

At the indicated times, mice were sacrificed and successive 15  $\mu$ m cryostat sections were prepared from the TA muscle. 1 every 5 slides was X-gal stained. Nuclei were counter-stained with Hoechst. The X-gal stained sections, were scored for the number of  $\beta$ -gal<sup>+</sup> nuclei and the obtained values were multiplied x 5 (total No. of nuclei). In addition  $\beta$ -gal<sup>+</sup>/total nuclei in 200  $\mu^2$  areas was counted. Results are reported in Table 2. Note that the increase in the  $\beta$ -gal<sup>+</sup>/total nuclei ratio is due to the

- 14 -

decrease in mononucleated cells which progressively occurs during muscle regeneration (fusion).

Table 2

No. of $\beta$ -gal <sup>+</sup> nuclei/muscle (0% of injected cells)		No. of $\beta$ -gal <sup>+</sup> nuclei/injection site (200 $\mu^2$ ) (0% of donor nuclei)	
Time(weeks)	MyoD fibroblasts	satellite cells	MyoD fibroblasts
0	1.0x10 <sup>6</sup> (100)	1.0x10 <sup>6</sup> (100)	
1	2.1x10 <sup>3</sup> (0.2)	0.5x10 <sup>3</sup> (0.05)	8/242 (3.3)
2	1.8x10 <sup>3</sup> (0.18)	1.9x10 <sup>3</sup> (0.19)	7/127 (5.5)
4	1.6x10 <sup>3</sup> (0.16)	3.3x10 <sup>3</sup> (0.33)	6/99 (6.0)
8	1.9x10 <sup>3</sup> (0.19)	3.6x10 <sup>3</sup> (0.36)	7/56 (12.5)
			2/256 (0.8)
			6/123 (4.8)
			8/105 (8.5)
			12/60 (13.1)

**Example 7**

**Muscle regeneration by genetically-modified fibroblasts expressing MyoD**

LIBSON

Fibroblasts isolated from human fetal skin were expanded and  
5 transduced in vitro with a high-titer stock of the retroviral vector  
carrying a lacZ gene encoding a cytoplasmic form of  $\beta$ -galactosidase  
(Salvatori, G., et al., 1993).

Transduction efficiency, estimated by the number of  $\beta$ -gal-staining  
cells in culture, ranged between 40 and 70%, making further selection of  
10 the transduced cells (e.g., G418 resistance) unnecessary. Transduced  
fibroblasts were exposed to the adeno-MyoD vector at an m.o.i. of 2,000,  
and then injected into the regenerating TA muscles of scid/bg mice (106  
cells/muscle in a single injection).

The efficiency of myogenic conversion of transduced fibroblasts  
15 was checked by allowing part of the cell culture to differentiate in vitro.  
On average, about 70% of the fibroblasts underwent myogenic conversion  
in these conditions, and most myotubes (> 90 %) scored positive for  
cytoplasmic  $\beta$ -gal expression. After 1, 2 and 4 weeks, mice were  
sacrificed and the TA serially sectioned and tibialis anterior muscles were  
20 cut and stained for  $\beta$ -gal activity.

Two weeks after injection,  $\beta$ -gal+ fibers were observed in 7 out of  
8 injected muscles. Higher magnification clearly revealed fibers  
accumulating the reporter gene product at variable levels, suggesting a  
variability in the proportion of injected/host cells in the  $\beta$ -gal+ fibers.

25 The presence and contribution of human cells in the newly formed  
fibers was confirmed by immunocytochemical analysis using an antibody  
which recognizes human but not mouse fetal myosin heavy chains, which

showed patches of intensive expression of human muscle-specific proteins in the regenerating areas.

In general, the average number of  $\beta$ -gal+ fibers per muscle obtained by injecting human converted fibroblasts was lower than that observed after injection of murine fibroblasts, although comparable to the number of positive cells observed in experiments in which lacZ-transduced human satellite cells were used. This indicates that human cells are, as expected, less efficient than murine ones in colonizing a mouse muscle.

However, and more importantly, human converted fibroblasts perform like primary myogenic cells in vivo, in terms of both number and size of newly formed fibers.

In order to better quantify the survival of human myogenically converted fibroblasts in mouse muscle, the Applicant also performed dot-blot Southern analysis with an Alu probe, as previously described (Salvatori, G., et al., 1993). To this purpose DNA was extracted from TA muscles of scid/bg mice two weeks after the injection of MyoD-converted human fibroblasts (following the same experimental protocol described above). As a positive control, the same number of human satellite cells was injected in similarly treated contra-lateral TA muscles.

The results of the dot-blot indicated that approximately 0.1% of the DNA extracted from TA injected with converted fibroblasts was of human origin; on the other hand this value raised to 0.3% in the case of the sample injected with human satellite cells.

Because histochemical analysis had revealed that virtually all the  $\beta$ -galactosidase staining was inside muscle fibers, we can conclude that the observed values faithfully reflect the percentage of human nuclei

incorporated into regenerated muscle fibers.

**Example 8**

**Immune response against cells exposed ex vivo to the adenoviral vector.**

5        One of the major problems associated to the in vivo use of  
adenoviral vectors in immunocompetent recipients is the induction of a  
significant, T-cell-mediated immune response against both the viral  
proteins and the product of the transgene(s). In the model described  
herein, no viral particle is injected directly in vivo. However, injected  
10    cells had been exposed shortly before administration to the adenoviral  
vector in vitro.

Replication-defective adenoviral DNA is expected to be rapidly  
diluted, and eventually lost, in cells in active cell division. Expression of  
the MyoD transgene, however, blocks almost immediately cell division in  
15    the converted fibroblasts, thus preventing the possibility of diluting the  
adenoviral vector by continuous culture.

To test whether an immune response could be raised by cells  
expressing a defective adenoviral genome in vivo, the Applicant injected  
2 x 10<sup>6</sup> C3H murine skin fibroblasts converted to myogenesis by  
20    exposure to the adeno-MyoD vector in the regenerating muscle of  
syngeneic mice.

Cells were labelled with DiI before injection and not with a lacZ  
transgene to avoid a possible immune reaction against the  $\beta$ -galactosidase  
protein. Animals were sacrificed 7, 14 and 21 days after the injection, and  
25    muscles analysed for the presence of immune infiltrate by  
immunofluorescence using antibodies against cell surface markers of  
murine leukocytes and macrophages.

No significant immune infiltrate was detected for up to 3 weeks after injection around the labelled cells, even though by this time all treated mice had developed antibodies against Ad5 proteins, which recognise adeno-infected but not control C3H fibroblasts

5        These experiments indicate that fibroblasts converted in vitro by exposure to an adeno-MyoD vector, and administered in vivo by intramuscular injection, do not elicit, and are therefore unlikely to be eliminated by, a cell-mediated cytotoxic immune response.

10       In contrast, direct injection of Adenoviral vector into regenerating muscle induced a strong immune infiltrate, as already reported (Yang, Y., et al., 1996).



## **BIBLIOGRAPHY**

- Salvatori, G., G. Ferrari, A. Mezzogiorno, S. Serivdei, M. Coletta, P. Tonali, R. Giavazzi, G. Cossu, and F. Mavilio. 1993. Retroviral vector-mediated gene transfer into human primary myogenic cells leads to expression in muscle fibers in vivo. *Human Gene Ther.* 4:713-723.
- Webster, C. e, H.M. Blau 1990. Accelerated age-related decline in replicative life span of Duchenne muscular dystrophy myoblasts: implication for cell and gene therapy. *Somatic Cell Mol. Genet.* 16:557-565.
- Davis, R. L., H. Weintraub, and A.B. Lassar. 1987. Expression of a single transfected cDNA converts fibroblasts to myoblasts. *Cell* 51:987-1000.
- Cossu, G., S. Tajbakhsh, and M. Buckingham. 1996. Myogenic specification in mammals. *Trends Genet.* 12:218-223.
- Cossu, G. 1997. Unorthodox myogenesis: possible developmental significance and implications for tissue histogenesis and regeneration. *Histol. Histopathol.* 12:755-760.
- Breton, M., Z., Li, D. Paulin, J.A. Harris, F. Rieger, M. Pincon-Raymond, and L. Garcia. 1995. Myotube driven myogenic recruitment of cells during in vitro myogenesis. *Develop. Dynam.* 202:126-136.
- Gibson, A.J., J. Karasinski, J. Relvas, J. Moss, T.G. Sherratt, P.N. Strong, and D.J. Watt. 1995. Dermal fibroblasts convert to a myogenic lineage in mdx mouse muscle. *J. Cell Science* 108:207-214.
- Salvatori, G., L. Lattanzi, M. Coletta, S. Aguanno, E. Vivarelli, R. Kelly, G. Ferrari, J. A. Harris, F. Mavilio, M. Molinaro, and G. Cossu. 1995. Myogenic conversion of mammalian fibroblasts induced by differentiating muscle cells. *J. Cell Science* 108:2733-2739.

Murry, C. E., M. A. Kay, T. Bartosek, S. D. Hauschka, and S. M. Schwartz. 1996. Muscle differentiation during repair of myocardial necrosis in rats via gene transfer with MyoD. *J. Clin. Invest.* 98:2209-2217.

5 Cusella - De Angelis, M.G., S. Molinari, A. Ledonne, M. Coletta, E. Vivarelli, M. Bouchè, M. Molinaro, S. Ferrari, and G. Cossu, 1994. Differential response of embryonic and fetal myoblasts to TGF $\beta$ : a possible regulatory mechanism of skeletal muscle histogenesis. *Development* 120:925-933.

10 Bader, D., T. Masaki, and D. A. Fischman. 1982. Immunochemical analysis of myosin heavy chain during avian myogenesis in vivo and in vitro. *J. Cell Biol.* 95:763-770.

Hasty, P., A. Bradley, J. H. Morris, D. G. Edmondson, J. M. Venuti, E. N. Olson, and W. H. Klein. 1993. Muscle deficiency and  
15 neonatal death in mice with a targeted mutation in the myogenin gene. *Nature* 375:787-790.

Edom, F., V. Mouly, J. P. Babert, M. Y. Fiszman, and G. S. Butler-Browne. 1994. Clones of human satellite cells can express in vitro both fast and slow myosin heavy chains. *Develop. Biology* 164:219-229.

20 Yasin, R., K. C. Van Beers, C. E. Nurse, S. Al-Ani, D. N. Landon, and E. J. Thompson. 1977. A quantitative technique for growing human adult skeletal muscle in culture starting from mononucleated cells. *J. Neurol. Sci.* 32:347-360.

Chomczynski, P., and N. Sacchi. 1987. Single-step method of RNA  
25 isolation by acid guanidinium thiocyanate-phenol-chlorophorm extraction. *Anal. Biochem.* 162:156-159.

Bober, E., G.E. Lyons, T. Braun, G. Cossu, M. Buckingham, and

H.H. Arnold. 1991. The muscle regulatory gene Myf-6 has a biphasic pattern of expression during early mouse development. *J. Cell Biol.* 113:1255-1265.

Boulter, J., W. Luyten, K. Evans, P. Mason, M. Ballivet, D. Goldman, D. Stengelin, S. Martin, S. Heinemann, and J. Patric, 1985. Isolation of a clone coding for the  $\alpha$ -subunit of a mouse acetylcholine receptor. *J. Neurosci.* 5:2545-2552.

Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*. 2nd ed., Cold Spring Harbor Laboratory Press, New York.

Yang, Y., S. E. Haecker, Q. Su, and J. Wilson, 1996. Immunology of gene therapy with adenoviral vectors. *Hum. Mol. Genet.* 5:1703-1712.

Tajbakhrsh, S., E. Vivarelli, M. G. Cusella-De Angelis, D. Rocancourt, M. Buckingham, and G. Cossu 1994. A population of myogenic cells derived from the mouse neural tube. *Neuron* 13:813-821.

Lyons, G. E., S. Muhlebach, A. Moser, R., Masood, B. M. Paterson, M. Buckingham, and J. C. Perriard. 1991. Development regulation of creatin kinase gene expression by myogenic factors in embryonic mouse and chick skeletal muscle. *Development* 113:1017-1029.

## CLAIMS

1. A method for the preparation of genetically-modified fibroblasts, expressing a "lineage commitment" gene, which comprises:
  - a) ex-vivo transduction of fibroblasts with a therapeutic gene or a gene capable of correcting a gene defect.
  - b) transient expression of the muscle lineage commitment gene in fibroblasts transduced as at point (a), through transformation of the cells with a high-efficiency vector, wherein the muscle lineage commitment gene is under the control of a strong promoter.
2. A method according to claim 1, wherein the therapeutic gene is chosen among the genes codifying sarcoglycanes, emerin, dystrophin, coagulation factors, insulin, growth hormone, hypoxanthine-guanine phosphoribosyl transferase, adenosine diaminase, purine nucleoside phosphorylase, glucocerebrosidase, low density lipoprotein receptor, phenylalanine hydroxylase, arginine succinate synthetase and aryl-sulfatase.
3. A method according to claim 2, wherein the therapeutical gene is the dystrophin gene.
4. A method according to claim 1, wherein the high efficiency vector is a non-viral vector selected from liposomes, polyethyleneimminas and nucleoprotein conjugates.
5. A method according to claim 1, wherein the high efficiency vector is a viral vector.
6. A method according to claim 5, wherein said viral vector is selected from baculovirus, adeno-related virus, adeno-virus.
7. A method according to claim 5, wherein said vector is adenovirus.
8. A method according to claim 1, wherein the muscle lineage commitment is selected from Myo-D, Myf-5, MRF4 and myogenin.
9. A method according to claim 8, wherein said gene is Myo-D.
10. A method according to claim 1, wherein the muscle lineage commitment gene is under the control of a viral promoter.
11. Genetically-modified fibroblasts transiently expressing a muscle lineage commitment gene.
12. Fibroblasts according to claim 11, wherein the muscle commitment lineage is Myo-D.

13. Fibroblasts according to claim 11 and 12, obtained with the method of claims 1-3.

Milan, May 8, 1998

The Mandatary

(Minoja Fabrizio)

of Bianchetti • Bracco • Minoja S.r.l.

(signature)